

Contribution of matter fields to the Gell-Mann-Low function for N=1 supersymmetric Yang-Mills theory, regularized by higher covariant derivatives.

K.V.Stepanyantz

February 1, 2008

*Moscow State University, physical faculty,
department of theoretical physics.
119992, Moscow, Russia*

Abstract

Contribution of matter fields to the Gell-Mann-Low function for N=1 supersymmetric Yang-Mills theory, regularized by higher covariant derivatives, is obtained using Schwinger-Dyson equations and Slavnov-Taylor identities. A possible deviation of the result from the corresponding contribution in the exact Novikov, Shifman, Vainshtein and Zakharov β -function is discussed.

1 Introduction.

It is well known, that supersymmetry essentially improves the ultraviolet behavior of a theory. For example, even in theories with unextended supersymmetry, it is possible to propose the form of the β -function exactly to all orders of the perturbation theory. This proposal was first made in Ref. [1] as a result of investigating instanton contributions structure. For the $N = 1$ supersymmetric Yang-Mills theory with matter fields, this β -function, called the exact Novikov-Shifman-Vainshtein-Zakharov (NSVZ) β -function) is

$$\beta(\alpha) = -\frac{\alpha^2 \left[3C_2 - 2C(R)(1 - \gamma(\alpha)) \right]}{2\pi(1 - C_2\alpha/2\pi)}, \quad (1)$$

where $\gamma(\alpha)$ is the anomalous dimension of the matter superfield. We note that here we present the result for a theory with the Dirac fermions, and two chiral superfields correspond to each of them. R denotes a representation for one of the matter superfields, and $C(R)$ is given by

$$\text{tr}(T^a T^b) = C(R) \delta^{ab}. \quad (2)$$

In order to obtain such β -function it is most convenient to use the higher derivative regularization [2, 3]. Then the renormalization of the operator $W_a C^{ab} W_b$ is exhausted at the one-loop, while the Gell-Mann-Low function coincides with the exact NSVZ β -function. This was confirmed by the explicit calculations in two-loop [4, 5], three-loop [6], and partially four-loop approximations [7]. (We note that the Gell-Mann-Low function does not depend on a particular choice of the renormalization scheme due to its definition.)

Investigation of the $N = 1$ supersymmetric electrodynamics was made in Ref. [8] exactly to all orders of the perturbation theory. According to that paper, it is possible to obtain the exact β -function using Schwinger-Dyson equations and Ward identities. However, there is an additional contribution besides the ordinary NSVZ result. Explicit calculations show that it is always equal to 0 in the lowest orders. This allows proposing existence of a new identity for some Green functions. A proof of this identity exactly to all orders is rather nontrivial. The matter is that it is not reduced to the Ward identity, while its proving by summation of diagrams [9] is rather complicated. We note that in Ref. [9] this proof was made only for a restricted class of diagrams in the massless case. Nevertheless, it seems that the new identity is also valid for the other diagrams, that follows, for example, from the explicit four-loop calculations [7].

So, obtaining the exact β -function is a rather nontrivial problem even in the electrodynamics. In the case of the Yang-Mills theory the exact β -function agrees with the results of explicit calculations in the two-loop approximation if the dimensional reduction is used for regularization. However, there are disagreements between the predictions, following from the exact β -function, and explicit calculations, made in $\overline{\text{MS}}$ -scheme for the β -function, defined as a derivative of the renormalized coupling constant, in the three-loop approximation [10, 11, 12]. It was shown [11] that these disagreements can be eliminated by a special choice of renormalization scheme, the possibility of such a choice being highly nontrivial [13].

In this paper we attempt to investigate contribution of the matter superfields to the Gell-Mann-Low function in non-Abelian supersymmetric gauge theories exactly to all orders of the perturbation theory using the Schwinger-Dyson equations and Slavnov-Taylor identities. The Gell-Mann-Low function is defined by asymptotic of the two-point Green function in the limit of large momentums. And so, the obtained results actually correspond to the limit of massless matter fields. (Nevertheless, in order to elucidate calculating the contribution of Pauli-Villars fields, we will consider the case of an arbitrary mass throughout the paper. Taking the limit $m \rightarrow 0$ will be made in the end for obtaining the final result.)

This paper is organized as follows.

In Sec. 2 we recall basic information about the $N = 1$ supersymmetric Yang-Mills theory and its higher derivatives regularization. The Schwinger-Dyson equations for the considered theory are obtained in Sec. 3. In that section we present the solution of the Slavnov-Taylor identities, which determines vertex functions in the Schwinger-Dyson equations. After substitution of these vertexes the contribution to the two-point Green function of the gauge field is obtained in the form of an integral. Calculation of this integral is made in Sec. 4. A brief discussion of the results is given in the conclusion.

2 $N = 1$ supersymmetric Yang-Mills theory, background field method and higher derivative regularization

The $N = 1$ supersymmetric Yang-Mills theory in the superspace is described by the action

$$S = \frac{1}{4e^2} \text{Re tr} \int d^4x d^2\theta W_a C^{ab} W_b + \frac{1}{4} \int d^4x d^4\theta \left(\phi^+ e^{2V} \phi + \tilde{\phi}^+ e^{-2V^t} \tilde{\phi} \right) + \frac{1}{2} m \int d^4x d^2\theta \tilde{\phi}^t \phi + \frac{1}{2} m \int d^4x d^2\bar{\theta} \tilde{\phi}^+ \phi^*. \quad (3)$$

Here ϕ and $\tilde{\phi}$ are chiral matter superfields, and V is a real scalar superfield, which contains the gauge field A_μ as a component. The superfield W_a is a supersymmetric analogue of the gauge field stress tensor. It is defined by

$$W_a = \frac{1}{32} \bar{D}(1 - \gamma_5) D \left[e^{-2V} (1 + \gamma_5) D_a e^{2V} \right], \quad (4)$$

where

$$D = \frac{\partial}{\partial \theta} - i\gamma^\mu \theta \partial_\mu \quad (5)$$

is a supersymmetric covariant derivative. In our notation, the gauge superfield V is expanded over the generators of the gauge group T^a as $V = e V^a T^a$, where e is a coupling constant.

Action (3) is invariant under gauge transformations

$$\phi \rightarrow e^{i\Lambda} \phi; \quad \tilde{\phi} \rightarrow e^{-i\Lambda^t} \tilde{\phi}; \quad e^{2V} \rightarrow e^{i\Lambda^+} e^{2V} e^{-i\Lambda}. \quad (6)$$

Such a transformation law means that if the field ϕ is in the representation R of the gauge group G , the field $\tilde{\phi}$ is in the representation $\bar{R}$, conjugated to R .

For quantization of this model it is convenient to use the background field method. The matter is that the background field method allows calculating the effective action without manifest breaking of the gauge invariance. In the supersymmetric case it can be formulated as follows [14, 15]: Let us make a substitution

$$e^{2V} \rightarrow e^{2V'} \equiv e^{\mathbf{\Omega}^+} e^{2V} e^{\mathbf{\Omega}}, \quad (7)$$

in action (3), where $\mathbf{\Omega}$ is a background scalar superfield. Expression for V' is a complicated nonlinear function of V , $\mathbf{\Omega}$, and $\mathbf{\Omega}^+$. We do not interest in explicit form of this function:

$$V' = V'[V, \mathbf{\Omega}]. \quad (8)$$

(For brevity of notation we will not explicitly write the dependence on $\mathbf{\Omega}^+$ here and below.)

The obtained theory will be invariant under background gauge transformations

$$\begin{aligned}\phi &\rightarrow e^\Lambda \phi; & \tilde{\phi} &\rightarrow e^{-\Lambda^t} \tilde{\phi}; & V &\rightarrow e^{iK} V e^{-iK}; \\ e^\Omega &\rightarrow e^{iK} e^\Omega e^{-\Lambda}; & e^{\Omega^+} &\rightarrow e^{-\Lambda^+} e^{\Omega^+} e^{-iK},\end{aligned}\tag{9}$$

where K is a real superfield, and Λ is a chiral superfield.

Let us construct the chiral covariant derivatives

$$\mathbf{D} \equiv e^{-\Omega^+} \frac{1}{2} (1 + \gamma_5) D e^{\Omega^+}; \quad \bar{\mathbf{D}} \equiv e^\Omega \frac{1}{2} (1 - \gamma_5) D e^{-\Omega}.\tag{10}$$

Acting on some field X , which is transformed as $X \rightarrow e^{iK} X$, these covariant derivatives are transformed in the same way. It is also possible to define a covariant derivative with the Lorentz index

$$\mathbf{D}_\mu \equiv -\frac{i}{4} (C\gamma^\mu)^{ab} \{ \mathbf{D}_a, \bar{\mathbf{D}}_b \},\tag{11}$$

which will have the same property. It is easy to see that after substitution (7) action (3) will be

$$\begin{aligned}S = & \frac{1}{4e^2} \text{tr Re} \int d^4x d^2\theta \mathbf{W}^a \mathbf{W}_a - \frac{1}{128e^2} \text{tr Re} \int d^4x d^4\theta \left[16 \left(e^{-2V} \mathbf{D}^a e^{2V} \right) \mathbf{W}_a + \right. \\ & + \left(e^{-2V} \mathbf{D}^a e^{2V} \right) \bar{\mathbf{D}}^2 \left(e^{-2V} \mathbf{D}_a e^{2V} \right) \Big] + \frac{1}{4} \int d^4x d^4\theta \left(\phi^+ e^{\Omega^+} e^{2V} e^\Omega \phi + \tilde{\phi}^+ e^{-\Omega^*} e^{-2V^t} \times \right. \\ & \left. \times e^{-\Omega^t} \tilde{\phi} \right) + \frac{1}{2} m \int d^4x d^2\theta \tilde{\phi}^t \phi + \frac{1}{2} m \int d^4x d^2\bar{\theta} \tilde{\phi}^+ \phi^*,\end{aligned}\tag{12}$$

where

$$\mathbf{W}_a = \frac{1}{32} e^\Omega \bar{\mathbf{D}} (1 - \gamma_5) D \left(e^{-\Omega} e^{-\Omega^+} (1 + \gamma_5) D_a e^{\Omega^+} e^\Omega \right) e^{-\Omega},\tag{13}$$

and the notation

$$\begin{aligned}\mathbf{D}^2 &\equiv \frac{1}{2} \bar{\mathbf{D}} (1 + \gamma_5) \mathbf{D}; & \bar{\mathbf{D}}^2 &\equiv \frac{1}{2} \bar{\mathbf{D}} (1 - \gamma_5) \mathbf{D}; \\ \mathbf{D}^a &\equiv \left[\frac{1}{2} \bar{\mathbf{D}} (1 + \gamma_5) \right]^a; & \mathbf{D}_a &\equiv \left[\frac{1}{2} (1 + \gamma_5) \mathbf{D} \right]_a; \\ \bar{\mathbf{D}}^a &\equiv \left[\frac{1}{2} \bar{\mathbf{D}} (1 - \gamma_5) \right]^a; & \bar{\mathbf{D}}_a &\equiv \left[\frac{1}{2} (1 - \gamma_5) \mathbf{D} \right]_a\end{aligned}\tag{14}$$

is used. Action of the covariant derivatives on the field V in the adjoint representation is defined by the standard way.

It is convenient to choose a regularization and gauge fixing so that invariance (9) will be unbroken. For example, the higher covariant derivative regularization can be introduced by the substitution of action (12) for

$$\begin{aligned}
S_\Lambda = & \frac{1}{4e^2} \text{tr Re} \int d^4x d^2\theta \mathbf{W}^a \mathbf{W}_a - \frac{1}{128e^2} \text{tr Re} \int d^4x d^4\theta \left[16 \left(e^{-2V} \mathbf{D}^a e^{2V} \right) \mathbf{W}_a + \right. \\
& + \left(e^{-2V} \mathbf{D}^a e^{2V} \right) \bar{\mathbf{D}}^2 \left(e^{-2V} \mathbf{D}_a \left(1 + \frac{(\mathbf{D}_\mu^2)^n}{\Lambda^{2n}} \right) e^{2V} \right) \Big] + \frac{1}{4} \int d^4x d^4\theta \left(\phi^+ e^{\Omega^+} e^{2V} e^{\Omega} \phi + \right. \\
& \left. + \tilde{\phi}^+ e^{-\Omega^*} e^{-2V} e^{-\Omega^t} \tilde{\phi} \right) + \frac{1}{2} m \int d^4x d^2\theta \tilde{\phi}^t \phi + \frac{1}{2} m \int d^4x d^2\bar{\theta} \tilde{\phi}^+ \phi^*. \tag{15}
\end{aligned}$$

This action is evidently invariant under background gauge transformations (9).

We note that the described way of regularization is a bit different from the method, proposed in Ref. [16]. The difference is in the form of the term, which contains higher derivatives. Using of the method, proposed here, simplifies calculations in a certain degree, while all particular features of higher derivative regularization are the same in the both cases. So, although explicit calculations by the method, proposed in Ref. [16] in two and more loops were not yet made, it is possible to propose that their results will not essentially differ from the ones, obtained below in this paper.

The gauge can be also fixed by the invariant way, for example, by adding the terms

$$S_{gf} = -\frac{1}{64e^2} \int d^4x d^4\theta \left(V \mathbf{D}^2 \bar{\mathbf{D}}^2 \left(1 + \frac{(\mathbf{D}_\mu^2)^n}{\Lambda^{2n}} \right) V + V \bar{\mathbf{D}}^2 \mathbf{D}^2 \left(1 + \frac{(\mathbf{D}_\mu^2)^n}{\Lambda^{2n}} \right) V \right). \tag{16}$$

In this case terms quadratic in the superfield V will have the simplest form:

$$\frac{1}{4e^2} \text{tr Re} \int d^4x d^4\theta V \mathbf{D}_\mu^2 \left(1 + \frac{(\mathbf{D}_\mu^2)^n}{\Lambda^{2n}} \right) V. \tag{17}$$

Certainly, it is also necessary to add the corresponding action S_{gh} for the Faddeev-Popov ghosts. We will not write here explicit form of this action, because it is not essential for calculating a contribution of the matter superfield loops.

Proposed way of the regularization and gauge fixing preserves both invariance under the supersymmetry transformations and the invariance under transformations (9). As a consequence, the effective action, calculated by the background field method, will be invariant under both supersymmetry and gauge transformations.

Let us construct the generating functional as follows:

$$\begin{aligned}
Z[J, \Omega, j] = & \int D\mu \exp \left\{ iS_\Lambda^r + iS_{gf} + iS_{gh} + iS_S + iS_{\phi_0} + \right. \\
& \left. + i \int d^4x d^4\theta \left(J + J[\Omega] \right) \left(V'[V, \Omega] - \mathbf{V} \right) \right\}, \tag{18}
\end{aligned}$$

where the superfield $\mathbf{V}$ is given by

$$e^{2\mathbf{V}} \equiv e^{\Omega^+} e^{\Omega}, \tag{19}$$

and $J[\Omega]$ is a so far undefined functional. A reason of its introducing will be clear later. The functional integration measure is written as

$$D\mu = DV D\phi D\tilde{\phi}. \quad (20)$$

S_Λ^r denotes renormalized and regularized action

$$\begin{aligned} S_\Lambda^r = & \frac{1}{4e_0^2} \text{tr Re} \int d^4x d^2\theta \mathbf{W}^a \mathbf{W}_a - \frac{1}{128e_0^2} \text{tr Re} \int d^4x d^4\theta \left[16 \left(e^{-2V} \mathbf{D}^a e^{2V} \right) \mathbf{W}_a + \right. \\ & + \left(e^{-2V} \mathbf{D}^a e^{2V} \right) \bar{\mathbf{D}}^2 \left(e^{-2V} \mathbf{D}_a \left(1 + \frac{(\mathbf{D}_\mu^2)^n}{\Lambda^{2n}} \right) e^{2V} \right) \Big] + \frac{1}{4} Z(e, \Lambda/\mu) \int d^4x d^4\theta \left(\phi^+ e^{\mathbf{\Omega}^+} \times \right. \\ & \times e^{2V} e^{\mathbf{\Omega}} \phi + \tilde{\phi}^+ e^{-\mathbf{\Omega}^*} e^{-2V^t} e^{-\mathbf{\Omega}^t} \tilde{\phi} \Big) + \frac{1}{2} m \int d^4x d^2\theta \tilde{\phi}^t \phi + \frac{1}{2} m \int d^4x d^2\bar{\theta} \tilde{\phi}^+ \phi^*, \end{aligned} \quad (21)$$

where e_0 is a bare coupling constant, and Z is a renormalization constant for the wave function of the matter superfields. S_{gf} denotes gauge fixing terms (16) (the coupling constant e in them is also replaced by e_0), and S_{gh} is a corresponding action for the Faddeev-Popov ghosts (and also for the Nielsen-Kallosh ghosts). S_S denotes the terms with sources for chiral superfields, which in the extended form are written as

$$S_S = \int d^4x d^2\theta \left(j^t \phi + \tilde{j}^t \tilde{\phi} \right) + \int d^4x d^2\bar{\theta} \left(j^+ \phi^* + \tilde{j}^+ \tilde{\phi}^* \right). \quad (22)$$

Moreover, in generating functional (18) we introduce the supplementary sources

$$\begin{aligned} S_{\phi_0} = & \frac{1}{4} \int d^4x d^4\theta \left(\phi_0^+ e^{\mathbf{\Omega}^+} e^{2V} e^{\mathbf{\Omega}} \phi + \phi^+ e^{\mathbf{\Omega}^+} e^{2V} e^{\mathbf{\Omega}} \phi_0 + \right. \\ & \left. + \tilde{\phi}_0^+ e^{-\mathbf{\Omega}^*} e^{-2V^t} e^{-\mathbf{\Omega}^t} \tilde{\phi} + \tilde{\phi}^+ e^{-\mathbf{\Omega}^*} e^{-2V^t} e^{-\mathbf{\Omega}^t} \tilde{\phi}_0 \right), \end{aligned} \quad (23)$$

where ϕ_0 , ϕ_0^+ , $\tilde{\phi}_0$ and $\tilde{\phi}_0^+$ are arbitrary scalar superfields. In principle, it is not necessary to introduce the term S_{ϕ_0} in the generating functional, but the presence of the parameters ϕ_0 is highly desirable for investigating the Schwinger-Dyson equations.

In order to understand how generating functional (18) is related with the ordinary effective action, we perform the substitution $V \rightarrow V'$. Then we obtain

$$Z[J, \mathbf{\Omega}, j] = \exp \left\{ -i \int d^4x d^4\theta \left(J + J[\mathbf{\Omega}] \right) \mathbf{V} \right\} Z_0 \left[J + J[\mathbf{\Omega}], \mathbf{\Omega}, j \right], \quad (24)$$

where

$$Z_0[J, \mathbf{\Omega}, j] = \int D\mu \exp \left\{ iS_\Lambda^r + iS_{gf} + iS_{gh} + iS_S + iS_{\phi_0} + i \int d^4x d^4\theta JV \right\}. \quad (25)$$

If the dependence of S_Λ^r , S_{gf} , S_{gh} , and S_{ϕ_0} on the arguments V , $\mathbf{\Omega}$, and $\mathbf{\Omega}^+$ were factorized into the dependence on the variable V' , Z_0 would not depend on $\mathbf{\Omega}$ and $\mathbf{\Omega}^+$ and would coincide with the ordinary generating functional. This really takes place for action (3) and S_{ϕ_0} . However, in the term with the higher derivatives and in the gauge fixing terms such factorization does not occur. Therefore, Z_0 actually differs from the ordinary generating functional.

Using the functional $Z[J, \mathbf{\Omega}, j]$ it is possible to construct the generating functional for the connected Green functions

$$\begin{aligned} W[J, \mathbf{\Omega}, j] &= -i \ln Z[J, \mathbf{\Omega}, j] = \\ &= - \int d^4x d^4\theta \left(J + J[\mathbf{\Omega}] \right) \mathbf{V} + W_0 \left[J + J[\mathbf{\Omega}], \mathbf{\Omega}, j \right]. \end{aligned} \quad (26)$$

Also it is possible to construct the corresponding effective action

$$\begin{aligned} \Gamma[V, \mathbf{\Omega}, \phi] &= - \int d^4x d^4\theta \left(J\mathbf{V} + J[\mathbf{\Omega}]\mathbf{V} \right) + W_0 \left[J + J[\mathbf{\Omega}], \mathbf{\Omega}, j \right] - \\ &- \int d^4x d^4\theta J V - \int d^4x d^2\theta \left(j^t \phi + \tilde{j}^t \tilde{\phi} \right) - \int d^4x d^2\bar{\theta} \left(j^+ \phi^* + \tilde{j}^+ \tilde{\phi}^* \right), \end{aligned} \quad (27)$$

where the sources should be expressed in terms of fields using the equation

$$\begin{aligned} V &= \frac{\delta}{\delta J} W[J, \mathbf{\Omega}, j] = -\mathbf{V} + \frac{\delta}{\delta J} W_0 \left[J + J[\mathbf{\Omega}], \mathbf{\Omega}, j \right]; \\ \phi &= \frac{\delta}{\delta j} W[J, \mathbf{\Omega}, j] = \frac{\delta}{\delta j} W_0 \left[J + J[\mathbf{\Omega}], \mathbf{\Omega}, j \right] \quad \text{e.t.c.} \end{aligned} \quad (28)$$

Substituting these expression into Eq. (27), we write the effective action as

$$\begin{aligned} \Gamma[V, \mathbf{\Omega}, \phi] &= W_0 \left[J + J[\mathbf{\Omega}], \mathbf{\Omega}, j \right] - \int d^4x d^4\theta \left(J[\mathbf{\Omega}]\mathbf{V} + J \frac{\delta}{\delta J} W_0 \left[j, J + J[\mathbf{\Omega}], \mathbf{\Omega}, j \right] \right) - \\ &- \int d^4x d^2\theta \phi \frac{\delta}{\delta j} W_0 \left[J + J[\mathbf{\Omega}], \mathbf{\Omega}, j \right] - \left(\text{similar terms with } \phi^+, \tilde{\phi} \text{ and } \tilde{\phi}^+ \right). \end{aligned} \quad (29)$$

Let us now set $V = 0$, so that

$$\mathbf{V} = \frac{\delta}{\delta J} W_0 \left[J + J[\mathbf{\Omega}], \mathbf{\Omega}, j \right]. \quad (30)$$

Moreover, we take into account that the invariance under background gauge transformations (9) essentially restricts the form of the effective action. If the quantum field V in the effective action is set to 0, the superfield K will affect only the gauge transformation law of the fields $\mathbf{\Omega}$ and $\mathbf{\Omega}^+$, and the only invariant combination is expression (19). (It is invariant in a sense, that the corresponding transformation law does not contain the superfield K .) This means that in the final expression for the effective action we can set

$$\mathbf{\Omega} = \mathbf{\Omega}^+ = \mathbf{V}. \quad (31)$$

In this case the effective action is

$$\begin{aligned} \Gamma[0, \mathbf{V}, \phi] &= W_0 \left[J + J[\mathbf{V}], \mathbf{V}, j \right] - \int d^4x d^4\theta \left(J + J[\mathbf{V}] \right) \frac{\delta}{\delta J} W_0 \left[J + J[\mathbf{V}], \mathbf{V}, j \right] - \\ &- \int d^4x d^2\theta \phi \frac{\delta}{\delta j} W_0 \left[J + J[\mathbf{V}], \mathbf{V}, j \right] - \left(\text{similar terms with } \phi^+, \tilde{\phi}, \text{ and } \tilde{\phi}^+ \right). \end{aligned} \quad (32)$$

Note, that this expression does not depend on form of the functional $J[\Omega]$. In particular it can be chosen to cancel terms linear in the field V in Eq. (18). Such a choice will be very convenient below.

If the gauge fixing terms and the terms with higher derivatives depended only on V' , expression (32) would coincide with the ordinary effective action. However, as we already mentioned above, the dependence on V , Ω , and Ω^+ is not factorized into the dependence on V' in the proposed method of renormalization and gauge fixing. According to Ref. [17, 18] the invariant charge (and, therefore, the Gell-Mann-Low function) is gauge independent, and the dependence of the effective action on gauge can be eliminated by renormalization of the wave functions of the gauge field, ghosts, and matter fields. Therefore, for calculating the Gell-Mann-Low function we may use the background gauge described above. We note that if this gauge is used, the renormalization constant of the gauge field A_μ is 1 due to the invariance of the action under transformations (9).

Nevertheless, generating functional (18) is not yet completely constructed. The matter is that adding the term with higher derivatives does not remove divergences from one-loop diagrams. To regularize them, it is necessary to insert the Pauli-Villars determinants in the generating functional [19]. Because in this paper we are interested only in contributions of matter superfields, we construct these determinants only for them. (Also it is necessary to introduce the Pauli-Villars fields for the gauge field and ghosts.) Here we will at once use condition (31). So, we insert in the generating functional the factor

$$\prod_i \left(\det PV(V, \mathbf{V}, M_i) \right)^{c_i}, \quad (33)$$

in which the Pauli-Villars determinants are defined by

$$\left(\det PV(\mathbf{V}, M) \right)^{-1} = \int D\Phi D\tilde{\Phi} \exp \left(iS_{PV} \right). \quad (34)$$

If condition (31) is used, the action for the Pauli-Villars fields will be

$$\begin{aligned} S_{PV} \equiv & Z(e, \Lambda/\mu) \frac{1}{4} \int d^4x d^4\theta \left(\Phi^+ e^{\mathbf{V}} e^{2V} e^{\mathbf{V}} \Phi + \tilde{\Phi}^+ e^{-\mathbf{V}^t} e^{-2V^t} e^{-\mathbf{V}^t} \tilde{\Phi} \right) + \\ & + \frac{1}{2} \int d^4x d^2\theta M \tilde{\Phi}^t \Phi + \frac{1}{2} \int d^4x d^2\bar{\theta} M \tilde{\Phi}^+ \Phi^*. \end{aligned} \quad (35)$$

The coefficients c_i in Eq. (33) satisfy conditions

$$\sum_i c_i = 1; \quad \sum_i c_i M_i^2 = 0. \quad (36)$$

Below, we assume that $M_i = a_i \Lambda$, where a_i are some constants. Inserting the Pauli-Villars determinants allows cancelling the remaining divergences in all one-loop diagrams, including diagrams containing counterterm insertions.

Making the transformation

$$\phi \rightarrow \phi/\sqrt{Z}; \quad \tilde{\phi} \rightarrow \tilde{\phi}/\sqrt{Z}; \quad \Phi \rightarrow \Phi/\sqrt{Z}; \quad \tilde{\Phi} \rightarrow \tilde{\Phi}/\sqrt{Z} \quad (37)$$

in the generating functional, we obtain that the dependence on the renormalization constant Z can be easily found from the generating functional, which does not contain the constant Z . If ϕ_0 and j are 0, this is made by the substitution

$$m \rightarrow m/Z; \quad M_i \rightarrow M_i/Z. \quad (38)$$

Therefore, we will make calculations with $Z = 1$, and dependence on the renormalization constant will be restored in the final result.

3 Schwinger-Dyson equations and Slavnov-Taylor identities.

In order to construct the Schwinger-Dyson equations for the considered theory it is necessary to split the action into three parts: the action for the background field, the kinetic term for quantum fields, which does not contain the background field, and interaction, in which the other terms are included:

$$S = S(\mathbf{V}) + S_2(V, \phi) + S_I(V, \mathbf{V}, \phi). \quad (39)$$

(Earlier we saw that the terms of the first order in the superfield V , which were obtained from the expansion of the classical action, can be omitted.) So, generating functional (18) can be written as

$$\begin{aligned} Z[J, j, \mathbf{V}] &= \int d\mu \exp \left(iS[\mathbf{V}] + iS_2[V, \phi] + iS_I[V, \mathbf{V}, \phi] + iS_S + i \int d^8x JV \right) = \quad (40) \\ &= \exp \left(iS[\mathbf{V}] + iS_I \left[\frac{1}{i} \frac{\delta}{\delta J}, \frac{1}{i} \frac{\delta}{\delta \mathbf{J}}, \frac{1}{i} \frac{\delta}{\delta j} \right] \right) \times \\ &\quad \times \int d\mu \exp \left(iS_2[V, \phi] + iS_S + i \int d^8x JV + i \int d^8x \mathbf{J}\mathbf{V} \right) \Big|_{\mathbf{J}=0}, \end{aligned}$$

where $\int d^8x \equiv \int d^4x d^4\theta_x$. Let us differentiate this expression with respect to the background field

$$\begin{aligned} \frac{\delta}{\delta \mathbf{V}_x} Z[J, j, \mathbf{V}] &= i \frac{\delta S[\mathbf{V}]}{\delta \mathbf{V}} Z + \exp \left(iS_I \left[\frac{1}{i} \frac{\delta}{\delta J}, \frac{1}{i} \frac{\delta}{\delta \mathbf{J}}, \frac{1}{i} \frac{\delta}{\delta j} \right] \right) i\mathbf{J}_x \times \\ &\quad \times \int d\mu \exp \left(iS_2[V, \phi] + iS_S + i \int d^8x JV + i \int d^8x \mathbf{J}\mathbf{V} \right) \Big|_{\mathbf{J}=0}. \quad (41) \end{aligned}$$

Moving the current $\mathbf{J}_x$ to the left and dividing the result to Z , we obtain

$$\frac{\delta}{\delta \mathbf{V}_x} W[J, j, \mathbf{V}] = \frac{\delta S[\mathbf{V}]}{\delta \mathbf{V}_x} + \frac{\delta}{\delta \mathbf{V}_x} S_I \left[\mathbf{V}, V + \frac{1}{i} \frac{\delta}{\delta J}, \phi + \frac{1}{i} \frac{\delta}{\delta j} \right] \quad (42)$$

Because the background field $\mathbf{V}$ is a parameter of the effective action, these equality can be equivalently written as

$$\frac{\delta \Gamma}{\delta \mathbf{V}_x} = \frac{\delta S[\mathbf{V}]}{\delta \mathbf{V}_x} + \left\langle \frac{\delta}{\delta \mathbf{V}_x} S_I \left[\mathbf{V}, V, \phi \right] \right\rangle \quad (43)$$

where the angular brackets denote taking an expectation value by the ordinary functional integration. Certainly, it is necessary to set the field V (an argument of the effective action) to 0 in the final result in accordance with Eq. (30) and the corresponding discussion.

Let us find a contribution to this expression given by the matter superfields. The corresponding interaction terms are

$$S_I = \frac{1}{4} \int d^8x \phi^+ \left(e^{\mathbf{V}} e^{2V} e^{\mathbf{V}} - 1 \right) \phi + \text{similar terms with } \tilde{\phi}. \quad (44)$$

Differentiating S_I with respect to the background field, we obtain that the corresponding contribution to the effective action is written in the form

$$\begin{aligned} \frac{\delta \Gamma}{\delta \mathbf{V}_x^a} = & \text{terms without matter} + \sum_i c_i \frac{\delta}{\delta \mathbf{V}_x^a} \left\langle \ln \det PV(V, \mathbf{V}, M_i) \right\rangle + \\ & + \frac{1}{4} \frac{\delta}{\delta \mathbf{V}_x^a} \int d^8x \left\langle \phi_x^+ e^{\mathbf{V}_x} e^{2V_x} e^{\mathbf{V}_x} \phi_x + \text{similar terms with } \tilde{\phi} \right\rangle. \end{aligned} \quad (45)$$

We are interested in the two-point Green function of the gauge field, corresponding to the expansion of the effective action in powers of the background field up to the second order terms. To calculate such expansion, it is convenient to use the following equality

$$\begin{aligned} \int d^8x d^8y \mathbf{V}_x^a \mathbf{V}_y^b \frac{\delta^2}{\delta \mathbf{V}_x^a \delta \mathbf{V}_y^b} \left(e^{\mathbf{V}} e^{2V} e^{\mathbf{V}} \right) \Big|_{\mathbf{V}=0} &= \mathbf{V} \left(\mathbf{V} e^{2V} + e^{2V} \mathbf{V} \right) + \left(\mathbf{V} e^{2V} + e^{2V} \mathbf{V} \right) \mathbf{V} = \\ &= e \int d^8x d^8y \mathbf{V}_y^b \mathbf{V}_x^a \left[T^a \frac{\delta}{\delta \mathbf{V}_y^b} \left(e^{\mathbf{V}} e^{2V} e^{\mathbf{V}} \right) \Big|_{\mathbf{V}=0} + \frac{\delta}{\delta \mathbf{V}_y^b} \left(e^{\mathbf{V}} e^{2V} e^{\mathbf{V}} \right) \Big|_{\mathbf{V}=0} T^a \right], \end{aligned} \quad (46)$$

where e in the beginning of the second string is a coupling constant. Using the form of the action for supplementary sources (23), we easily obtain

$$\frac{1}{4} \left\langle \phi_x^+ T^a e^{\mathbf{V}_x} e^{2V_x} e^{\mathbf{V}_x} \phi_x + \phi_x^+ e^{\mathbf{V}_x} e^{2V_x} e^{\mathbf{V}_x} T^a \phi_x \right\rangle = \frac{1}{i} \text{tr} \left[T^a \left(\frac{\delta^2 \Gamma}{\delta j_x^+ \delta \phi_{0x}^+} + \frac{\delta^2 \Gamma}{\delta j_x \delta \phi_{0x}} \right) \right]. \quad (47)$$

(In the last equation the matrix notation is used for the brevity.) The derivatives with respect to the sources must be expressed in terms of fields as follows:

$$\begin{aligned} \frac{\delta}{\delta j_x^+} = & \int d^8z \left[\left(\frac{\delta^2 \Gamma}{\delta \phi_z \delta \phi_x^+} \right)^{-1} \frac{D^2}{8\partial^2} \frac{\delta}{\delta \phi_z} + \left(\frac{\delta^2 \Gamma}{\delta \phi_z^+ \delta \phi_x^+} \right)^{-1} \frac{\bar{D}^2}{8\partial^2} \frac{\delta}{\delta \phi_z^+} + \right. \\ & \left. + \left(\frac{\delta^2 \Gamma}{\delta \tilde{\phi}_z \delta \phi_x^+} \right)^{-1} \frac{D^2}{8\partial^2} \frac{\delta}{\delta \tilde{\phi}_z} + \left(\frac{\delta^2 \Gamma}{\delta \tilde{\phi}_z^+ \delta \phi_x^+} \right)^{-1} \frac{\bar{D}^2}{8\partial^2} \frac{\delta}{\delta \tilde{\phi}_z^+} + \left(\frac{\delta^2 \Gamma}{\delta V_z \delta \phi_x^+} \right)^{-1} \frac{\delta}{\delta V_z} \right]. \end{aligned} \quad (48)$$

Therefore, using Eqs. (45), (46), and taking into account similar terms with the fields $\tilde{\phi}$, the corresponding contribution to the two-point Green function of the matter superfield can be written as

$$\frac{\delta^2 \Gamma}{\delta \mathbf{V}_y^b \delta \mathbf{V}_x^a} = \dots + e \frac{\delta}{\delta \mathbf{V}_y^b} \text{tr} \left[T^a \frac{1}{i} \frac{\delta^2 \Gamma}{\delta j_x^+ \delta \phi_{0x}^+} - (T^a)^t \frac{1}{i} \frac{\delta^2 \Gamma}{\delta \tilde{j}_x^+ \delta \tilde{\phi}_{0x}^+} + \text{h.c.} \right], \quad (49)$$

where dots denote contributions of the gauge fields, ghosts, and also all possible Pauli-Villars fields. (In this expression we omit symmetrization with respect to the indexes a and b , because, as we will see below, it will be symmetric automatically.) We note that the calculation of the Pauli-Villars fields contributions are made completely similar to the calculation of ordinary fields contributions [8], and details of this calculation are not presented here. The result will be given below.

In order to calculate expressions in Eq. (49), it is necessary to use Eq. (48) and manifestly perform the differentiation. Then the result can be graphically presented as a sum of two effective diagrams

$$\Delta \Gamma_V^{(2)} = \text{diagram 1} + \text{diagram 2} \quad (50)$$

The double lines correspond to the effective propagators, which are written as

$$\left(\frac{\delta^2 \Gamma}{\delta \phi_x^+ \delta \phi_y} \right)^{-1} = - \frac{G D_x^2 \bar{D}_x^2}{4(\partial^2 G^2 + m^2 J^2)} \delta_{xy}^8; \quad \left(\frac{\delta^2 \Gamma}{\delta \phi_x \delta \tilde{\phi}_y} \right)^{-1} = - \frac{m J \bar{D}_x^2}{\partial^2 G^2 + m^2 J^2} \delta_{xy}^8, \quad (51)$$

depending on the chirality of the ends. The functions G and J are determined by the two-point Green functions of the matter superfield as

$$\frac{\delta^2 \Gamma}{\delta \phi_x^+ \delta \phi_y} = \frac{D_x^2 \bar{D}_x^2}{16} G(\partial^2) \delta_{xy}^8; \quad \frac{\delta^2 \Gamma}{\delta \phi_x \delta \tilde{\phi}_y} = - \frac{\bar{D}_x^2}{4} m J(\partial^2) \delta_{xy}^8, \quad (52)$$

where $\delta_{xy}^8 \equiv \delta^4(x-y) \delta^4(\theta_x - \theta_y)$, and the subscripts denote points where the expressions are evaluated.

The vertex functions can be obtained from the Slavnov-Taylor identities similar to the case of the electrodynamics. Certainly, it is necessary to take into account that we consider the vertex functions, which have an external line of the **background** field. As we already mentioned above, the effective action is invariant under the background gauge transformations. It is easy to see that this invariance can be expressed by the equality

$$0 = \int d^8 y \left(\frac{\delta \Gamma}{\delta \mathbf{V}_y^a} \delta \mathbf{V}_y^a + \frac{\delta \Gamma}{\delta \phi_y} \frac{D^2}{8 \partial^2} \delta \phi_y + \delta \phi_y^+ \frac{\bar{D}^2}{8 \partial^2} \frac{\delta \Gamma}{\delta \phi_y^+} + \frac{\delta \Gamma}{\delta \phi_{0y}} \delta \phi_{0y} + \delta \phi_{0y}^+ \frac{\delta \Gamma}{\delta \phi_{0y}^+} + \right. \\ \left. + \text{similar terms with } \tilde{\phi} \text{ and } \tilde{\phi}_0 \right), \quad (53)$$

where

$$\delta \phi_x = \Lambda_x \phi_x; \quad \delta \phi_x^+ = \phi_x^+ \Lambda_x^+; \quad \delta \phi_{0x} = \Lambda_x \phi_{0x}; \quad \delta \phi_{0x}^+ = \phi_{0x}^+ \Lambda_x^+; \\ \delta \mathbf{V}_x = - \frac{1}{2} (\Lambda_x + \Lambda_x^+) + O(\mathbf{V}). \quad (54)$$

Here Λ is an arbitrary chiral superfield, and all other terms in $\delta\mathbf{V}$ are proportional at least to the first degree of the background field. (For the fields $\tilde{\phi}$ e.t.c. the transformation laws can be also easily written.) Let us differentiate Eq. (53) with respect to Λ_y^a , ψ_z^+ , ϕ_x or with respect to Λ_y^a , $\tilde{\psi}_z$, ϕ_x . As a result we obtain the Slavnov-Taylor identities

$$\begin{aligned} 0 &= \frac{1}{2e}(\bar{D}_y^2 + D_y^2) \frac{\delta^3\Gamma}{\delta\mathbf{V}_y^a \delta\phi_{0z}^+ \delta\phi_x} - \frac{\delta^2\Gamma}{\delta\phi_{0z}^+ \delta\phi_y} T^a \bar{D}_y^2 \delta_{xy}^8 - D_y^2 \left(\delta_{yz}^8 T^a \frac{\delta^2\Gamma}{\delta\phi_{0y}^+ \delta\phi_x} \right); \\ 0 &= \frac{1}{2e}(\bar{D}_y^2 + \bar{D}_y^2) \frac{\delta^3\Gamma}{\delta\mathbf{V}_y^a \delta\tilde{\phi}_{0z} \delta\phi_x} - \frac{\delta^2\Gamma}{\delta\tilde{\phi}_{0z} \delta\phi_y} T^a \bar{D}_y^2 \delta_{xy}^8 + \bar{D}_y^2 \left(\delta_{yz}^8 T^a \frac{\delta^2\Gamma}{\delta\tilde{\phi}_{0y} \delta\phi_x} \right). \end{aligned} \quad (55)$$

where

$$\frac{\delta^2\Gamma}{\delta\phi_y \delta\phi_{0z}^+} = -\frac{1}{8}G(\partial^2)\bar{D}_y^2 \delta_{yz}^8; \quad \frac{\delta^2\Gamma}{\delta\tilde{\phi}_y^* \delta\phi_{0z}^+} = \frac{m}{32\partial^2} \left(J(\partial^2) - 1 \right) D_y^2 \bar{D}_y^2 \delta_{yz}^8, \quad (56)$$

They differ from the similar identities for the electrodynamics only in the presence of gauge group generators. Therefore, the solution will be also similar

$$\begin{aligned} \left. \frac{\delta^3\Gamma}{\delta\mathbf{V}_y^a \delta\phi_{0z}^+ \delta\phi_x} \right|_{p=0} &= e \left[-2\partial^2 \Pi_{1/2y} \left(\bar{D}_y^2 \delta_{xy}^8 \delta_{yz}^8 \right) F(q^2) + \frac{1}{8} D^b C_{bc} \bar{D}_y^2 \left(\bar{D}_y^2 \delta_{xy}^8 D_y^c \delta_{yz}^8 \right) f(q^2) - \right. \\ &\quad \left. - \frac{1}{16} q^\mu G'(q^2) \bar{D} \gamma^\mu \gamma_5 D_y \left(\bar{D}_y^2 \delta_{xy}^8 \delta_{yz}^8 \right) - \frac{1}{4} \bar{D}_y^2 \delta_{xy}^8 \delta_{yz}^8 G(q^2) \right] T^a; \end{aligned} \quad (57)$$

$$\begin{aligned} \left. \frac{\delta^3\Gamma}{\delta\mathbf{V}_y^a \delta\tilde{\phi}_{0z} \delta\phi_x} \right|_{p=0} &= e \left[\frac{m}{32} D^b C_{bc} D_y^2 \left(D_y^2 \bar{D}_y^2 \delta_{xy}^8 D_y^c \delta_{yz}^8 \right) h(q^2) + \frac{m}{16} J'(q^2) \left(\bar{D}_y^2 \delta_{xy}^8 D_y^2 \delta_{yz}^8 - \right. \right. \\ &\quad \left. \left. - D_y^2 \bar{D}_y^2 \delta_{xy}^8 \delta_{yz}^8 \right) + \frac{m}{16} \left(\frac{J'(q^2)}{q^2} - \frac{J(q^2) - 1}{q^4} \right) \left(D_y^2 \bar{D}_y^2 \delta_{xy}^8 \frac{\bar{D}_y^2 D_y^2}{16} \delta_{yz}^8 + D_y^2 \bar{D}_y^2 \delta_{xy}^8 q^2 \delta_{yz}^8 \right) \right] T^a. \end{aligned} \quad (58)$$

The primes denote derivatives with respect to q^2 ,

$$\Pi_{1/2} = -\frac{1}{8\partial^2} D^a \bar{D}^2 D_a = -\frac{1}{8\partial^2} \bar{D}^a D^2 \bar{D}_a \quad (59)$$

is a supersymmetric transverse projection operator, notation for the derivatives are similar to (14), and the functions F , f and h can not be determined from the Slavnov-Taylor identities.

Functions (57) and (58) allow finding ordinary Green functions by the identities


$$-\frac{D_z^2}{2} \frac{\delta^3\Gamma}{\delta\mathbf{V}_x^a \delta\phi_y \delta\phi_{0z}^+} = \frac{\delta^3\Gamma}{\delta\mathbf{V}_x^a \delta\phi_y \delta\phi_z^+}; \quad -\frac{\bar{D}_z^2}{2} \frac{\delta^3\Gamma}{\delta\mathbf{V}_x^a \delta\phi_y \delta\tilde{\phi}_{0z}} = \frac{\delta^3\Gamma}{\delta\mathbf{V}_x^a \delta\phi_y \delta\tilde{\phi}_z} \quad (60)$$

We note that the supplementary sources in Eq. (23) were specially introduced in order that such identities take place. A proof of these relations can be made completely similar to the case of the electrodynamics, considered in [8] in details. Using Eqs. (60) we find

$$\left. \frac{\delta^3 \Gamma}{\delta \mathbf{V}_y^a \delta \phi_z^+ \delta \phi_x} \right|_{p=0} = e \left[\partial^2 \Pi_{1/2y} \left(\bar{D}_y^2 \delta_{xy}^8 D_y^2 \delta_{yz}^8 \right) F(q^2) + \right. \\ \left. + \frac{1}{32} q^\mu G'(q^2) \bar{D} \gamma^\mu \gamma_5 D_y \left(\bar{D}_y^2 \delta_{xy}^8 D_y^2 \delta_{yz}^8 \right) + \frac{1}{8} \bar{D}_y^2 \delta_{xy}^8 D_y^2 \delta_{yz}^8 G(q^2) \right] T^a; \quad (61)$$

$$\left. \frac{\delta^3 \Gamma}{\delta \mathbf{V}_y^a \delta \tilde{\phi}_z \delta \phi_x} \right|_{p=0} = -\frac{em}{32} J'(q^2) \left[\bar{D}_y^2 \delta_{xy}^8 D_y^2 \bar{D}_y^2 \delta_{yz}^8 - D_y^2 \bar{D}_y^2 \delta_{xy}^8 \bar{D}_y^2 \delta_{yz}^8 \right] T^a. \quad (62)$$

Obtained expressions for the vertex functions allow calculating all expressions, entering Eq. (49) in the limit of zero external momentum. For this purpose it is convenient to make the following observation: The first diagram in Eq. (50) can be equivalently presented in the form



$$+ e \frac{\delta}{\delta \mathbf{V}_y^b} \text{tr} \left[(T^a)^t \frac{m \bar{D}_x^2}{16 \partial^2} \left(\frac{\delta^2 \Gamma}{\delta \phi_z^+ \delta \tilde{\phi}_x^+} \right)^{-1}_{z=z} - T^a \frac{m \bar{D}_x^2}{16 \partial^2} \left(\frac{\delta^2 \Gamma}{\delta \tilde{\phi}_z^+ \delta \phi_x^+} \right)^{-1}_{z=z} + \text{h.c.} \right]. \quad (63)$$

The single line in the second diagram corresponds to

$$- \frac{D^2 \bar{D}^2}{16 \partial^2} \delta_{12}^8 \quad (64)$$

if fields at the ends 1 and 2 have opposite chirality, and 0, if chirality of the ends is the same. Eq. (63) can be easily verified using Feynman rules and performing the differentiation with respect to the field $\mathbf{V}$.

We will calculate the expression

$$\left. \frac{\partial}{\partial \ln \Lambda} \frac{\delta \Gamma}{\delta \mathbf{V}_y^b \delta \mathbf{V}_x^a} \right|_{p=0}. \quad (65)$$

We note that the regularization by higher covariant derivatives is essentially used here, because it allows differentiating the integrand and taking the limit of zero external momentum.

Using Eqs. (51), (61), and (62) it is easy to see that

$$e \frac{\partial}{\partial \ln \Lambda} \frac{\delta}{\delta \mathbf{V}_y^b} \text{tr} \left[(T^a)^t \frac{m \bar{D}_x^2}{16 \partial^2} \left(\frac{\delta^2 \Gamma}{\delta \phi_z^+ \delta \tilde{\phi}_x^+} \right)^{-1}_{z=z} - T^a \frac{m \bar{D}_x^2}{16 \partial^2} \left(\frac{\delta^2 \Gamma}{\delta \tilde{\phi}_z^+ \delta \phi_x^+} \right)^{-1}_{z=z} + \text{h.c.} \right]_{p=0} = \\ = e^2 C(R) \delta^{ab} \partial_{xy}^8 \frac{\partial}{\partial \ln \Lambda} \int \frac{d^4 q}{(2\pi)^4} \frac{1}{q^2} \frac{d}{dq^2} \left(\frac{m^2 J}{q^2 G^2 + m^2 J^2} \right). \quad (66)$$

(All similar expressions are written in the Euclidean space after the Weak rotation.)

Let us proceed to the calculation of all other contributions in Eq. (50). For this purpose we note that they contain the sum of subdiagrams, presented in Fig. 1. The




Figure 1: The sum of subdiagrams in the Schwinger-Dyson equations.

external line in these subdiagrams corresponds to the background field $\mathbf{V}$, and expressions for propagators were presented above. After simple transformations it is easy to see that expressions for the sum of these subdiagrams in the limit $p \rightarrow 0$ are written as

$$\begin{aligned}
& D_y^2 \bar{D}_y^2 \left(\mathbf{V}_y \frac{G D_y^2 \bar{D}_y^2}{64 \partial^2 (\partial^2 G^2 + m^2 J^2)} \delta_{yz}^8 \right) + \mathbf{V}_y \frac{G D_y^2 \bar{D}_y^2}{4 (\partial^2 G^2 + m^2 J^2)} \delta_{yz}^8 = \\
& = - \int d^8 x \frac{D_x^2}{16 \partial^2} \delta_{xy}^8 \left(2 \partial^2 \Pi_{1/2} \mathbf{V}_x + i \bar{D} \gamma^\mu \gamma_5 D \mathbf{V}_x \partial_\mu \right) \frac{G \bar{D}_z^2}{\partial^2 G^2 + m^2 J^2} \delta_{xz}^8 - \\
& - D^a \mathbf{V}_y \frac{G D_a \bar{D}_z^2}{2 (\partial^2 G^2 + m^2 J^2)} \delta_{yz}^8 - D^2 \mathbf{V}_y \frac{G \bar{D}_z^2}{4 (\partial^2 G^2 + m^2 J^2)} \delta_{yz}^8
\end{aligned} \tag{67}$$

or

$$\begin{aligned}
& = D_y^2 \bar{D}_y^2 \left(\mathbf{V}_y \frac{m J D_y^2}{16 \partial^2 (\partial^2 G^2 + m^2 J^2)} \delta_{yz}^8 \right) + \mathbf{V}_y \frac{m J D_y^2}{\partial^2 G^2 + m^2 J^2} \delta_{yz}^8 = \\
& = \int d^8 x \frac{D_x^2}{64 \partial^2} \delta_{xy}^8 \left(2 \partial^2 \Pi_{1/2} \mathbf{V}_x + i \bar{D} \gamma^\mu \gamma_5 D \mathbf{V}_x \partial_\mu \right) \frac{m J \bar{D}_x^2 D_x^2}{\partial^2 (\partial^2 G^2 + m^2 J^2)} \delta_{xz}^8 + \\
& + \frac{1}{8} D^a \mathbf{V}_y \frac{m J D_{ay} \bar{D}_y^2 D_y^2}{\partial^2 (\partial^2 G^2 + m^2 J^2)} \delta_{yz}^8 + \frac{1}{16} D_y^2 \mathbf{V}_y \frac{m J \bar{D}_y^2 D_y^2}{\partial^2 (\partial^2 G^2 + m^2 J^2)} \delta_{yz}^8
\end{aligned} \tag{68}$$

depending on the chirality of the ends. Terms in subdiagrams, containing the integral over $d^8 x$, have chiral projection operators, acting on the ends y and z . Due to identities (60), this allows using simpler vertexes (61) and (62) for the calculations. After simple, but rather long calculations we obtain that the contribution of these terms to Eq. (65) is

$$e^2 C(R) \delta^{ab} \partial^2 \Pi_{1/2} \delta_{xy}^8 \frac{\partial}{\partial \ln \Lambda} \int \frac{d^4 q}{(2\pi)^4} \frac{1}{q^2} \frac{d}{dq^2} \ln \left(q^2 G^2 + m^2 J^2 \right). \tag{69}$$

It is also easy to see that a contribution of the terms, which are proportional to $D^2 \mathbf{V}$ in Eqs. (67) and (68), is 0. However, a contribution of the terms, proportional to $D^a \mathbf{V}$ is nontrivial and contains the unknown functions f and h . It can be written as

$$\begin{aligned}
& \text{Diagram: A circle with a wavy line attached to a vertex on the left. The vertex is labeled } D^a V. \text{ The circle is labeled } I_a. \\
& = \frac{\partial}{\partial \ln \Lambda} \text{tr} \int d^8 x d^8 y \mathbf{V}_y D^a \mathbf{V}_x \frac{D_{az} \bar{D}_z^2}{4 i \partial^2} \frac{\delta^3 \Gamma}{\delta j_z^+ \delta \mathbf{V}_y \delta \phi_{0x}^+} \Big|_{z=x, p=0},
\end{aligned} \tag{70}$$

where the symbol I_a (we use the notation in [9] here) means that in the expression for the exact propagator it is necessary to make the substitution

$$\bar{D}^2 D^2 \delta_{xz}^8 \rightarrow 2 \bar{D}^2 D^a \delta_{xz}^8. \quad (71)$$

After substituting propagators and vertex functions the considered contribution will be

$$e^2 C(R) \delta^{ab} \partial^2 \Pi_{1/2} \delta_{xy}^8 \frac{\partial}{\partial \ln \Lambda} \int \frac{d^4 q}{(2\pi)^4} \left(- \frac{m^2 J (2q^2 J' - (J-1))}{q^4 (q^2 G^2 + m^2 J^2)} - \frac{16Gf + 16m^2 Jh}{q^2 G^2 + m^2 J^2} \right). \quad (72)$$

Collecting all contributions to the effective action, we obtain the following result ($Z = 1$, a dependence on Z will be restored later):

$$\begin{aligned} \left. \frac{\partial}{\partial \ln \Lambda} \frac{\delta^2 \Gamma}{\delta \mathbf{V}_y^b \delta \mathbf{V}_x^a} \right|_{p=0} = \dots + e^2 C(R) \delta^{ab} \partial^2 \Pi_{1/2} \delta_{xy}^8 \frac{\partial}{\partial \ln \Lambda} \int \frac{d^4 q}{(2\pi)^4} \left\{ \frac{1}{q^2} \frac{d}{dq^2} \left(\ln (q^2 G^2 + \right. \right. \\ \left. \left. + m^2 J^2) + \frac{m^2 J}{q^2 G^2 + m^2 J^2} \right) - \frac{m^2 J (2q^2 J' - (J-1))}{q^4 (q^2 G^2 + m^2 J^2)} - \frac{16Gf + 16m^2 Jh}{q^2 G^2 + m^2 J^2} - \right. \\ \left. - \text{similar terms for the Pauli-Villars fields} \right\}, \quad (73) \end{aligned}$$

where dots denote contributions of the gauge fields and ghosts. Comparing the obtained result with the corresponding expression for the supersymmetric electrodynamics, we see that they differ in the multiplier $C(R)$. Nevertheless, one more difference is possible: In the electrodynamics explicit calculations show that there is an identity, which can be written as equality to 0 of contribution (70). The problem, if a similar identity is valid in the non-Abelian case, is not yet solved. In general, it is not necessary, because there are new diagrams, which contain a matter loop and vertexes with self-action of the gauge superfield, in the non-Abelian case. That is why in this paper we will not assume existence of such identity.

4 Exact Gell-Mann-Low function.

Investigating the supersymmetric electrodynamics [8] reveals that there are divergences only in the one-loop approximation if the higher derivative regularization is used, while the Gell-Mann-Low function coincides with the exact NSVZ β -function. Let us consider a non-Abelian theory.

The Gell-Mann-Low function is defined by the dependence of finite part of the two-point Green function on the momentum in the limit $m \rightarrow 0$. And so, let us consider the massless case and write the renormalized two-point Green function as

$$\Gamma_V^{(2)} = - \frac{1}{16\pi} \text{tr} \int \frac{d^4 p}{(2\pi)^4} \mathbf{V}(-p) \partial^2 \Pi_{1/2} \mathbf{V}(p) d^{-1}(\alpha, \mu/p), \quad (74)$$

where α is a renormalized coupling constant. It is known [20, 21] that the function d is invariant under renormgroup transformations, i.e. does not depend on a choice of

the renormalization prescription. The Gell-Mann-Low function, which we will denote by $\beta(\alpha)$, is given by

$$\beta\left(d(\alpha, \mu/p)\right) = -\frac{\partial}{\partial \ln p} d(\alpha, \mu/p). \quad (75)$$

Because the function d is renorminvariant, the function $\beta(\alpha)$ will not also depend on a particular renormalization prescription. This is its difference from the function $b(\alpha)$, which is constructed from the renormalized coupling constant by the differentiation with respect to the logarithm of the normalization point.

In order to obtain physical results it is necessary to calculate the function d in the massive case, and then to impose the boundary condition

$$d(\alpha, \mu/p, m/\mu) \Big|_{p=0} = \alpha_{ph}, \quad (76)$$

where α_{ph} is an experimentally measured coupling constant, at a given μ . Moreover, it is necessary to impose one more boundary condition to the Green functions of the matter superfield:

$$G(\alpha, \mu/p, m/\mu) \Big|_{p^2=m_{ph}^2} = \frac{m}{m_{ph}} J(\alpha, \mu/p, m/\mu) \Big|_{p^2=m_{ph}^2}, \quad (77)$$

in which m_{ph} denotes a physical mass. (This equality assures that the propagator of the matter superfield will have a pole at the physical mass.) The boundary conditions written above allow expressing the parameters α and m (at a given μ) through the physically observed values. Nevertheless, the results of this paper do not allow finding explicit form of the functions d , G , and J in the massive case. And so, we can not relate α and m with the experimentally observed values. However, it is possible to obtain an expression for the contribution of matter superfields to the Gell-Mann-Low function:

If d_0 denotes the function d calculated at $Z = 1$, the following equation takes place in accordance with Eq. (73)

$$\begin{aligned} -\frac{\partial}{\partial \ln p} d_0^{-1}(\alpha_0, \Lambda/p) \Big|_{p=0} &= \frac{\partial}{\partial \ln \Lambda} d_0^{-1}(\alpha_0, \Lambda/p) \Big|_{p=0} = -16\pi C(R) \frac{\partial}{\partial \ln \Lambda} \times \\ &\times \int \frac{d^4 q}{(2\pi)^4} \frac{1}{2q^2} \frac{d}{dq^2} \left\{ \ln(q^2 G^2) + X(q^2/\Lambda^2) - \sum_i c_i \ln(q^2 G_{PV}^2 + M_i^2 J_{PV}^2) - \right. \\ &\left. - \sum_i c_i \frac{M_i^2 J_{PV}}{q^2 G_{PV}^2 + M_i^2 J_{PV}^2} - \sum_i c_i X(q^2/\Lambda^2, a_i) \right\}, \end{aligned} \quad (78)$$

where the dimensionless functions X are obtained from the equations

$$\begin{aligned} \frac{\partial X(q^2/\Lambda^2)}{\partial q^2} &= -\frac{16f}{G}; \\ \frac{\partial X(q^2/\Lambda^2, a_i)}{\partial q^2} &= -\frac{M_i^2 J (2q^2 J' - (J - 1))}{q^4 (q^2 G^2 + M_i^2 J^2)} - \frac{16Gf + 16M_i^2 Jh}{q^2 G^2 + M_i^2 J^2}, \end{aligned} \quad (79)$$

in which $M_i = a_i \Lambda$.

The obtained integral is reduced to the total derivative in the four-dimensional spherical coordinates, only the substitution at the low limit being different from 0 [8]:

$$\begin{aligned} \left. \frac{\partial}{\partial \ln \Lambda} d_0^{-1} \right|_{p=0} &= \frac{C(R)}{2\pi} \frac{\partial}{\partial \ln \Lambda} \left\{ \ln G(0)^2 + X(0) - \sum_i c_i \ln \left(M_i^2 J_{PV}(0, a_i)^2 \right) - \right. \\ &\quad \left. - \sum_i c_i \frac{1}{J_{PV}(0, a_i)} - \sum_i c_i X(0, a_i) \right\} = -\frac{1}{\pi} C(R) \left(1 - \frac{\partial \ln G}{\partial \ln \Lambda} - \frac{1}{2} \frac{\partial X}{\partial \ln \Lambda} \right) \Big|_{q=0} \end{aligned} \quad (80)$$

Here we took into account that the functions $X(q^2/\Lambda^2, a_i)$ and $J_{PV}(q^2/\Lambda^2, a_i)$ in the limit $q \rightarrow 0$ tended to some finite constants, because they were defined by convergent (even in the limit $\Lambda \rightarrow \infty$), dimensionless integrals, which did not contain infrared divergences. Because both parts of Eq. (80) depend on α_0 and Λ/p , it allows finding the expression for d_0^{-1} up to an insignificant numerical constant

$$d_0^{-1}(\alpha_0, \Lambda/p) = -\frac{1}{\pi} C(R) \left(\ln \frac{\Lambda}{p} - \ln G(\alpha_0, \Lambda/p) - \frac{1}{2} X(\alpha_0, \Lambda/p) \right) + \text{const.} \quad (81)$$

Making substitution (38) in Eq. (80) it is possible to restore the dependence of the effective action on the renormalization constant Z :

$$d^{-1}(\alpha, \mu/p) = d_0^{-1}(\alpha_0, \Lambda/p) + \frac{C(R)}{\pi} \ln Z(\alpha, \Lambda/\mu). \quad (82)$$

Differentiating this expression with respect to the momentum p , we obtain the Gell-Mann-Low function:

$$-\frac{\partial}{\partial \ln p} d^{-1} = -\frac{1}{\pi} C(R) \left(1 + \frac{\partial \ln G}{\partial \ln p} + \frac{1}{2} \frac{\partial X}{\partial \ln p} \right). \quad (83)$$

Using Eq. (79), we rewrite this contribution of the matter superfield as

$$-\frac{C(R)}{\pi} \left(1 - \gamma(\alpha) - \lim_{p \rightarrow 0} \frac{16p^2 f}{G} \right), \quad (84)$$

where $\gamma(\alpha)$ is the anomalous dimension of the matter superfield:

$$\gamma(d(\alpha, \mu/p)) = -\frac{\partial}{\partial \ln p} \ln ZG(\alpha, \mu/p). \quad (85)$$

(Expression (84) contributes to the Gell-Mann-Low function and should be compared with the numerator of Eq. (1).) Let us again mention that the obtained result does not depend on a choice of renormalization scheme.

We note that the contribution of the last term in Eq. (84) was always 0 in all diagrams calculated up to now. Moreover, if the identity

$$\text{tr} \frac{\partial}{\partial \ln \Lambda} \int d^8 x d^8 y \mathbf{V}_y D^a \mathbf{V}_x \frac{D_{az} \bar{D}_z^2}{4i \partial^2} \frac{\delta^3 \Gamma}{\delta j_z^+ \delta \mathbf{V}_y \delta \phi_{0x}^+} \Big|_{z=x, p=0} = 0, \quad (86)$$

is valid in the non-Abelian case, this will be always true. Nevertheless, in the non-Abelian case this statement requires an additional verification. That is why we for the generality write the correction, corresponding to a nontrivial contribution of the function f .

We note that the function f can actually affect only diagrams in three and more loops. Therefore, if such contribution appeared calculating the scheme dependent function $b(\alpha)$, it could be always eliminated by a proper redefinition of the coupling constant.

Finally we note that due to the similarity of the result with the electrodynamics case, if the contribution of the function f is absent, there are divergences only in the one-loop approximation. Details of the corresponding calculation are presented in Ref. [8]. This completely agrees with the structure of the anomalies supermultiplet and is a particular feature of higher covariant derivative regularization using [22].

5 Conclusion

In this paper a contribution of the matter superfields to the two-point Green function of the gauge field is calculated by the Schwinger-Dyson equations and Slavnov-Taylor identities for the supersymmetric Yang-Mills theory regularized by higher covariant derivatives. This regularization is very essential, because the method of calculating, used here, is based on the differentiation with respect to the regularization parameter and taking the limit of zero momentum.

In many respects the obtained results are similar to the corresponding results in the electrodynamics. However, it is necessary to mention some differences:

1. The calculation was made using the background field method, because this method allows preserving the manifest gauge invariance of the effective action. Without its using it would be also necessary to calculate the renormalization constant for the gauge superfield that make calculations essentially more complicated.

2. There is the multiplier $C(R)$, defined by Eq. (2) in the non-Abelian case. The same multiplier is also present in the exact NSVZ β -function.

3. Deviations of the result from the exact β -function, due to the contribution of the function f , which can not be determined from the Slavnov-Taylor identity, are possible hypothetically. Such contributions seem to disappear in the electrodynamics due to a new identity for the Green functions. However, there are diagrams, containing vertexes with self-action of the gauge field, in the non-Abelian case, which can break the additional identity. Contributions of these diagrams were not yet calculated explicitly in the considered regularization. That is why we also present the result for the β -function, if there is a nontrivial contribution of the function f .

We note that according to the above discussion, it is possible to point two possible reasons why the three-loop calculations with the dimensional reduction differ from the NSVZ β -function:

1. There is a nontrivial contribution of the function f .
2. In preceding papers instead of the Gell-Mann-Low function, the β -function, defined as a derivative of the renormalized coupling constant in the $\overline{\text{MS}}$ -scheme, is calculated.

At present we can not affirm which of this variants (or both ones simultaneously) is realized. This requires either explicit calculating or finding an origin of identity (86).

Moreover, it is necessary to calculate a contribution of the gauge fields and ghosts using Schwinger-Dyson equations and Slavnov-Taylor identities. Now this work is in progress.

Acknowledgments.

This paper was supported by the Russian Foundation for Basic Research (Grant No. 05-01-00541).

References

- [1] *V.Novikov, M.Shifman, A.Vainshtein, V.Zakharov*, Phys.Lett. **166B**, (1985), 329.
- [2] *A.A.Slavnov*, Theor.Math.Phys. **23**, (1975), 3.
- [3] *T.Bakeyev, A.Slavnov*, Mod.Phys.Lett. **A11**, (1996), 1539.
- [4] *A.Soloshenko, K.Stepanyantz*, Two-loop renormalization of $N = 1$ supersymmetric electrodynamics, regularized by higher derivatives, hep-th/0203118.
- [5] *A.Soloshenko, K.Stepanyantz*, Theor.Math.Phys., **134**, (2003), 377.
- [6] *A.Soloshenko, K.Stepanyantz*, Theor.Math.Phys. **140**, (2004), 1264.
- [7] *A.Pimenov, K.Stepanyantz*, Theor.Math.Phys., **147**, (2006), 687.
- [8] *K.V.Stepanyantz*, Theor.Math.Phys. **142**, (2005), 29.
- [9] *K.Stepanyantz*, Theor.Math.Phys. **146**, (2006), 321.
- [10] *L.V.Avdeev, O.V.Tarasov*, Phys.Lett. **112 B**, (1982), 356.
- [11] *I.Jack, D.R.T.Jones, C.G.North*, Nucl.Phys. **B 473**, (1996), 308
- [12] *I.Jack, D.R.T.Jones, C.G.North*, Phys.Lett. **386 B**, (1996), 138
- [13] *I.Jack, D.R.T.Jones*, Regularization of supersymmetric theories, hep-ph/ 9707278.
- [14] *P.West*, Introduction to supersymmetry and supergravity, World Scientific, 1986.
- [15] *J.Gates, M.Grisaru, M.Rocek, W.Siegel*, Front.Phys., **58**, (1983), 1.
- [16] *P.West*, Nucl.Phys. **B 268**, (1986), 113.
- [17] *H.Kluberg-Stern, J.B.Zuber*, Phys.Rev. **D12**, (1975), 467.
- [18] *H.Kluberg-Stern, J.B.Zuber*, Phys.Rev. **D12**, (1975), 482.
- [19] *L.D.Faddeev, A.A.Slavnov*, Gauge fields, introduction to quantum theory, second edition, Benjamin, Reading, 1990.

- [20] *N.N.Bogolyubov, D.V.Shirkov*, Introduction to the theory of quantized fields, Intersci.Monogr.Phys.Astron. **3**, (1959), 1.
- [21] *C.Itzykson, J.-B.Zuber*, Quantum field theory, McGraw-Hill Inc. 1980.
- [22] *K.Stepanyantz*, Theor.Math.Phys. **140**, (2004), 939.